



TOPIC : 5 - ODD AND EVEN FUNCTIONS

Even function:

If $f(x)$ is an even function, then

$$f(x) = f(-x) \Rightarrow b_n = 0.$$

Ex: $\cos x, |x|, x^2$
 $f(x) = x^2 \Rightarrow f(-x) = (-x)^2 = x^2 = f(x)$

$$\therefore f(x) = f(-x).$$

$\therefore f(x)$ is an even function $\Rightarrow b_n = 0.$

Odd function:

If $f(x)$ is an odd function, then

$$-f(x) = +f(-x).$$

Ex: $f(x) = x^3, \sin x, x^3, x \cos x$

$$\Rightarrow f(-x) = (-x)^3 = -x^3 = -f(x)$$

$$\therefore f(-x) = -f(x)$$

$\therefore f(x)$ is an odd function $\Rightarrow a_0 = 0$ & $a_n = 0$

Problems based on $(-\pi, \pi)$ & $(-l, l)$

First check whether the function is odd or even.

If the function is even using the fourier formula in the interval $(-\pi, \pi)$.

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos nx \quad \text{Here } b_n = 0.$$



If the function is odd using the fourier formula in the interval $(-\pi, \pi)$.

$$f(x) = \sum_{n=1}^{\infty} b_n \sin nx$$

$$\text{where } b_n = \frac{2}{b-a} \int_a^b f(x) \sin nx \, dx$$

Here $a_0 = 0$ & $a_n = 0$.

If the function is even using the fourier formula in the interval $(-l, l)$.

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos \frac{n\pi x}{l}$$

$$\text{where } a_0 = \frac{2}{b-a} \int_{-l}^l f(x) \, dx$$

$$a_n = \frac{2}{b-a} \int_a^b f(x) \cos \frac{n\pi x}{l} \, dx$$

If the function is odd using the fourier formula in the interval $(-l, l)$.

$$f(x) = \sum_{n=1}^{\infty} b_n \sin \frac{n\pi x}{l}$$

$$\text{where } b_n = \frac{2}{b-a} \int_{-l}^l f(x) \sin \frac{n\pi x}{l} \, dx$$



and deduce that

$$\frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots = \frac{\pi}{8}$$

Sol:

$$f(-x) = \begin{cases} 1 - \frac{2x}{\pi} & , \quad -\pi \leq -x \leq 0 \\ 1 + \frac{2x}{\pi} & , \quad 0 \leq -x \leq \pi \end{cases}$$

$$= \begin{cases} 1 - \frac{2x}{\pi} & , \quad 0 \leq x \leq \pi \\ 1 + \frac{2x}{\pi} & , \quad -\pi \leq x \leq 0 \end{cases}$$

$$f(-x) = f(x)$$

$\therefore f(x)$ is even $\Rightarrow b_n = 0$

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos nx$$

$$a_0 = \frac{2}{b-a} \int_a^b f(x) dx$$

$$= \frac{2}{2\pi} \int_{-\pi}^{\pi} f(x) dx$$

$$= \frac{2}{\pi} \int_0^{\pi} f(x) dx = \frac{2}{\pi} \int_0^{\pi} \left(1 - \frac{2x}{\pi}\right) dx$$

$$= \frac{2}{\pi} \left[x - \frac{2x^2}{2\pi} \right]_0^{\pi}$$

$$= \frac{2}{\pi} \left[\pi - \frac{\pi^2}{\pi} \right] = \frac{2}{\pi} \cdot 0$$

$$a_0 = 0$$

$$a_n = \frac{2}{b-a} \int_a^b f(x) \cos nx dx$$



$$a_n = \frac{2}{\pi} \int_0^{\pi} \left(1 - \frac{2x}{\pi}\right) \cos nx \, dx$$

Here $u = 1 - \frac{2x}{\pi}$ $\int dv = \int \cos nx \, dx$

$u_1 = -\frac{2}{\pi}$ $v = \frac{\sin nx}{n}$

$u_2 = 0$ $v_1 = -\frac{\cos nx}{n^2}$

$$a_n = \frac{2}{\pi} \left[\left(1 - \frac{2x}{\pi}\right) \frac{\sin nx}{n} - \frac{2 \cos nx}{\pi n^2} \right]_0^{\pi}$$

$$= \frac{2}{\pi} \left[\left(1 - \frac{2\pi}{\pi}\right) \frac{\sin n\pi}{n} - \frac{2 \cos n\pi}{\pi n^2} + \frac{2}{\pi n^2} \right]$$

$$= \frac{2}{\pi} \left[\frac{-2(-1)^n}{\pi n^2} + \frac{2}{\pi n^2} \right]$$

$$a_n = \frac{8}{\pi n^2} [1 - (-1)^n]$$

$$a_n = \begin{cases} \frac{8}{\pi n^2} & \text{if } n \text{ is odd} \\ 0 & \text{if } n \text{ is even} \end{cases}$$

$$f(x) = \sum_{n=\text{odd}}^{\infty} \frac{8}{\pi n^2} \cos nx$$

$$1 = \frac{8}{\pi} \sum_{n=\text{odd}}^{\infty} \frac{1}{n^2} \cos nx$$

$$\frac{\pi}{8} = \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots$$

$f(x) = |\cos x|$



and deduce that

$$\frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots = \frac{\pi}{8}$$

sol:

$$f(-x) = \begin{cases} 1 - \frac{2x}{\pi} & , \quad -\pi \leq -x \leq 0 \\ 1 + \frac{2x}{\pi} & , \quad 0 \leq -x \leq \pi \end{cases}$$

$$= \begin{cases} 1 - \frac{2x}{\pi} & , \quad 0 \leq x \leq \pi \\ 1 + \frac{2x}{\pi} & , \quad -\pi \leq x \leq 0 \end{cases}$$

$$f(-x) = f(x)$$

$\therefore f(x)$ is even $\Rightarrow b_n = 0$

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos nx$$

$$a_0 = \frac{2}{b-a} \int_a^b f(x) dx$$

$$= \frac{2}{2\pi} \int_{-\pi}^{\pi} f(x) dx$$

$$= \frac{2}{\pi} \int_0^{\pi} f(x) dx = \frac{2}{\pi} \int_0^{\pi} \left(1 - \frac{2x}{\pi}\right) dx$$

$$= \frac{2}{\pi} \left[x - \frac{2x^2}{2\pi} \right]_0^{\pi}$$

$$= \frac{2}{\pi} \left[\pi - \frac{\pi^2}{\pi} \right] = \frac{2}{\pi} \cdot \cancel{2\pi}$$

$$a_0 = 0$$

$$a_n = \frac{2}{b-a} \int_a^b f(x) \cos nx dx$$



$f(x)$ is even. $\Rightarrow b_n = 0$

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos nx.$$

$$a_0 = \frac{2}{b-a} \int f(x) dx$$

$$= \frac{2}{2\pi} \int_{-\pi}^{\pi} |\cos x| dx = \frac{2}{\pi} \int_0^{\pi} |\cos x| dx$$

$$= \frac{2}{\pi} \left\{ \int_0^{\pi/2} \cos x dx + \int_{\pi/2}^{\pi} (-\cos x) dx \right\}$$

$$= \frac{2}{\pi} \left\{ (\sin x) \Big|_0^{\pi/2} - (\sin x) \Big|_{\pi/2}^{\pi} \right\}$$

$$= \frac{2}{\pi} \left\{ \sin \frac{\pi}{2} - \sin 0 - \sin \pi + \sin \frac{\pi}{2} \right\}$$

$$= \frac{2}{\pi} \left\{ 2 \right\} = \frac{4}{\pi}$$

$$a_0 = \frac{4}{\pi}$$
$$a_n = \frac{2}{2\pi} \int_{-\pi}^{\pi} |\cos x| \cos nx dx$$

$$= \frac{2}{\pi} \int_0^{\pi} |\cos x| \cos nx dx$$

$$= \frac{2}{\pi} \left\{ \int_0^{\pi/2} \cos x \cos nx dx - \int_{\pi/2}^{\pi} \cos x \cos nx dx \right\}$$



$$\begin{aligned} a_n &= \frac{1}{\pi} \left[\left(\frac{\sin(n+1)x}{n+1} + \frac{\sin(n-1)x}{n-1} \right) \Big|_0^{\pi/2} \right. \\ &\quad \left. - \left(\frac{\sin(n+1)x}{n+1} + \frac{\sin(n-1)x}{n-1} \right) \Big|_{\pi/2}^{\pi} \right] \\ &= \frac{1}{\pi} \left[\frac{\sin(n+1)\pi/2}{n+1} + \frac{\sin(n-1)\pi/2}{n-1} + \frac{\sin(n+1)\pi/2}{n+1} \right. \\ &\quad \left. + \frac{\sin(n-1)\pi/2}{n-1} \right] \\ &= \frac{1}{\pi} \left[\frac{2 \sin(n+1)\pi/2}{n+1} + \frac{2 \sin(n-1)\pi/2}{n-1} \right] \\ &= \frac{2}{\pi} \left[\frac{\sin n\pi/2 \cos \pi/2 + \cos n\pi/2 \sin \pi/2}{n+1} \right. \\ &\quad \left. + \frac{\sin n\pi/2 \cos \pi/2 - \cos n\pi/2 \sin \pi/2}{n-1} \right] \\ &= \frac{2}{\pi} \left[\frac{\cos n\pi/2}{n+1} - \frac{\cos n\pi/2}{n-1} \right] \\ &= \frac{2}{\pi} \frac{\cos n\pi/2}{n^2-1} [n-1 - n-1] \\ a_n &= \frac{-4}{\pi} \frac{\cos n\pi/2}{n^2-1} \\ a_1 &= 2 \left\{ \int_0^{\pi/2} \cos x \cos x dx - \int_{\pi/2}^{\pi} \cos x \cos x dx \right\} \end{aligned}$$

$$= \frac{2}{\pi} \int_0^{\pi/2} \left(\frac{1 + \cos 2x}{2} \right) dx - \int_{\pi/2}^{\pi} \left(\frac{1 + \cos 2x}{2} \right) dx$$

$$a_1 = \frac{2}{\pi} \left[\left(\frac{1}{2}x + \frac{\sin 2x}{4} \right) \Big|_0^{\pi/2} - \left(\frac{1}{2}x + \frac{\sin 2x}{4} \right) \Big|_{\pi/2}^{\pi} \right]$$

$$= \frac{2}{\pi} \left[\frac{1}{2} \cdot \frac{\pi}{2} + \frac{\sin 2\pi/2}{4} - \frac{1}{2} \pi - \frac{\sin 2\pi}{4} + \frac{1}{2} \cdot \frac{\pi}{2} + \frac{\sin 2 \cdot \pi/2}{4} \right]$$

$$= \frac{2}{\pi} \left[\frac{\pi}{4} - \frac{\pi}{2} + \frac{\pi}{4} \right]$$

$$= \frac{2}{\pi} \left[\frac{2\pi}{4} - \frac{\pi}{2} \right]$$

$$a_1 = 0.$$

$$f(x) = \frac{4}{\pi \cdot 2} + \sum_{n=2}^{\infty} \frac{-4}{\pi} \cdot \frac{\cos n\pi/2}{n^2 - 1}$$

$$f(x) = \frac{2}{\pi} + \frac{4}{\pi} \sum_{n=2}^{\infty} \frac{\cos n\pi/2}{n^2 - 1}$$

Problems based on odd function:

1) Determine the Fourier series for the function

$$f(x) = \begin{cases} -1+x, & -\pi < x < 0 \\ 1+x, & 0 < x < \pi \end{cases} \quad \& \text{ hence deduce}$$

$$\text{Given } f(x) = \begin{cases} x-1 & -\pi < x < 0 \\ 1+x & 0 < x < \pi \end{cases}$$

$$f(-x) = \begin{cases} -x-1 & -\pi < -x < 0 \\ 1-x & 0 < -x < \pi \end{cases}$$

$$= \begin{cases} x+1 & 0 < x < \pi \\ x-1 & -\pi < x < 0 \end{cases}$$

$$f(-x) = -f(x)$$

$f(x)$ is an odd function $\Rightarrow a_0 = 0, a_n = 0$

$$\therefore f(x) = \sum_{n=1}^{\infty} b_n \sin nx$$

$$\text{where } b_n = \frac{2}{\pi} \int_0^{\pi} f(x) \sin nx \, dx$$

$$= \frac{2}{\pi} \int_0^{\pi} (x+1) \sin nx \, dx$$

$$= \frac{2}{\pi} \left[-(x+1) \frac{\cos nx}{n} + \frac{\sin nx}{n^2} \right]_0^{\pi}$$

$$= \frac{2}{\pi} \left[-(\pi+1) \frac{\cos n\pi}{n} + \frac{\sin n\pi}{n^2} + \frac{1}{n} \right]$$

$$= \frac{2}{\pi} \left[-(\pi+1) \frac{(-1)^n}{n} + \frac{1}{n} \right]$$

$$b_n = \frac{2}{n\pi} \left[1 - (\pi+1)(-1)^n \right]$$



2. Prove that $\frac{x(\pi^2 - x^2)}{12} = \frac{\sin x}{1^3} - \frac{\sin 2x}{2^3} + \frac{\sin 3x}{3^3} - \dots$
in the interval $(-\pi, \pi)$.

sol:

$$\text{Let } f(x) = \frac{x(\pi^2 - x^2)}{12}$$

$$f(-x) = (-x) \left(\frac{\pi^2 - (-x)^2}{12} \right)$$

$$= -x \left(\frac{\pi^2 - x^2}{12} \right)$$

$$f(-x) = -f(x)$$

$f(x)$ is an odd function $\Rightarrow a_0 = 0$ & $a_n = 0$

Let the required fourier series be

$$f(x) = \sum_{n=1}^{\infty} b_n \sin nx$$

$$b_n = \frac{2}{\pi} \int_0^{\pi} f(x) \sin nx \, dx$$

$$= \frac{2}{\pi} \int_0^{\pi} \frac{x(\pi^2 - x^2)}{12} \sin nx \, dx$$

$$u = x\pi^2 - x^3 \quad \int dv = \int \sin nx = \frac{2}{12\pi} \int_0^{\pi} (\pi^2 x^2 - x^3) \sin nx \, dx$$

$$u_1 = \pi^2 - 3x^2 \quad \downarrow \quad v_1 = \frac{-\cos nx}{n}$$

$$u_2 = -6x \quad \downarrow \quad v_2 = \frac{-\sin nx}{n^2}$$

$$u_3 = -6 \quad \downarrow \quad v_3 = \frac{\cos nx}{n^3}$$

$$= \frac{1}{6\pi} \left[(\pi^2 x - x^3) \frac{\cos nx}{n} + (\pi^2 - 3x^2) \frac{\sin nx}{n^2} \right.$$

$$\left. - 6x \frac{\cos nx}{n^2} + 6 \frac{\sin nx}{n^3} \right]_0^{\pi}$$

$$b_n = -\frac{\cos n\pi}{n^3}$$

$$= -\frac{(-1)^n}{n^3}$$

$$\therefore f(x) = -\sum_{n=1}^{\infty} \frac{(-1)^n}{n^3} \sin nx$$

$$\frac{x(\pi^2 - x^2)}{12} = \frac{\sin x}{1^3} - \frac{\sin 2x}{2^3} + \frac{\sin 3x}{3^3} + \dots$$

③ $f(x) = x^2 + x$ in $(-\pi, \pi)$ $\int_1^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6}$

Sol:

$$f(x) = x + x^2$$

$$f(-x) = -x + (-x)^2$$

$$= -x + x^2$$

$$f(x) \neq f(-x)$$

$$f(x) = -f(-x)$$

$\therefore f(x)$ is neither even nor odd.

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos nx + \sum_{n=1}^{\infty} b_n \sin nx$$

$$a_0 = \frac{2}{2\pi} \int_{-\pi}^{\pi} f(x) dx$$

$$= \frac{1}{\pi} \int_{-\pi}^{\pi} x + x^2 dx$$

$$= \frac{1}{\pi} \left(\frac{x^2}{2} + \frac{x^3}{3} \right)_{-\pi}^{\pi}$$

$$= \left[\frac{\pi^2}{2} + \frac{\pi^3}{3} - \frac{\pi^2}{2} + \frac{\pi^3}{3} \right]$$



$$\begin{aligned} a_n &= \frac{2}{2\pi} \int_{-\pi}^{\pi} f(x) \cos nx \, dx \\ &= \frac{1}{\pi} \int_{-\pi}^{\pi} x + x^2 \cos nx \, dx \\ &= \frac{1}{\pi} \left[(x + x^2) \frac{\sin nx}{n} + (1 + 2x) \frac{\cos nx}{n^2} - 2 \frac{\sin nx}{n^3} \right]_{-\pi}^{\pi} \\ &= \frac{1}{\pi} \left[(1 + 2\pi) \frac{\cos \pi n}{n^2} - (1 + 2(-\pi)) \frac{\cos n(-\pi)}{n^2} \right] \\ &= \frac{1}{\pi} \left[(1 + 2\pi) \frac{(-1)^n}{n^2} - (1 - 2\pi) \frac{(-1)^n}{n^2} \right] \\ &= \frac{(-1)^n}{n^2 \pi} [x + 2\pi - y + 2\pi] \\ &= \frac{(-1)^n 4\pi}{n^2 \pi} \end{aligned}$$

$$a_n = \frac{4(-1)^n}{n^2}$$

$$b_n = \frac{2}{2\pi} \int_{-\pi}^{\pi} x + x^2 \sin nx \, dx$$

$$= \frac{1}{\pi} \left[(x + x^2) \frac{\cos nx}{n} + (1 + 2x) \frac{\sin nx}{n^2} + \frac{2 \cos nx}{n^3} \right]_{-\pi}^{\pi}$$

$$= \frac{1}{\pi} \left[-(\pi + \pi^2) \frac{(-1)^n}{n} + \frac{2(-1)^n}{n^3} + \frac{(-\pi + \pi^2) \cos n(-\pi)}{n} - \frac{2 \cos n(-\pi)}{n^3} \right]$$



$$b_n = \frac{1}{\pi} \left[\frac{-\pi(-1)^n}{n} - \frac{\pi^2(-1)^n}{n} - \frac{\pi(-1)^n}{n} + \frac{\pi^2(-1)^n}{n} \right]$$

$$= \frac{1}{\pi} \left[\frac{-2\pi(-1)^n}{n} \right]$$

$$b_n = \frac{2(-1)^{n+1}}{n}$$

$$b_n = \begin{cases} \frac{2}{n} & \text{if } n \text{ is odd} \\ -\frac{2}{n} & \text{if } n \text{ is even} \end{cases}$$

$$f(x) = \frac{\pi^2}{3 \cdot 2} + \sum_{n=1}^{\infty} \frac{4(-1)^n}{n^2} \cos nx + \sum_{n=1}^{\infty} \frac{2(-1)^{n+1}}{n} \sin nx$$

If $x=0, x=\pi$

$$\pi + \pi = \frac{2\pi^2}{3} + \sum_{n=1}^{\infty} \frac{4(-1)^n}{n^2} \cos n\pi$$

$$\pi + \pi = \frac{2\pi^2}{3} = 4 \sum_{n=1}^{\infty} \frac{(-1)^n (-1)^n}{n^2}$$

If $x = \frac{\pi}{2}$ (discontinuous at end point)

$$\frac{f(-\pi) + f(\pi)}{2} = \frac{\pi^2}{3} + 4 \sum_{n=1}^{\infty} \frac{(-1)^n (-1)^n}{n^2}$$

$$\frac{\pi^2 - \pi^2}{2} = \frac{\pi^2}{3} + 4 \sum_{n=1}^{\infty} \frac{(-1)^{2n}}{n^2}$$

$$\pi^2 - \frac{\pi^2}{3} = 4 \sum_{n=1}^{\infty} \frac{(-1)^{2n}}{n^2}$$

$$\frac{2\pi^2}{3 \cdot 4} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \dots$$

$$\frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \dots = \frac{\pi^2}{6}$$